



ehponline.org

# ENVIRONMENTAL HEALTH PERSPECTIVES

## Early Life Exposure to Outdoor Air Pollution and Respiratory Health, Ear Infections, and Eczema in Infants from the INMA Study

**Inmaculada Aguilera, Marie Pedersen, Raquel Garcia-Esteban,  
Ferran Ballester, Mikel Basterrechea, Ana Esplugues,  
Ana Fernández-Somoano, Aitana Lertxundi,  
Adonina Tardón, and Jordi Sunyer**

**<http://dx.doi.org/10.1289/ehp.1205281>**

**Online 4 December 2012**



**NIEHS**

National Institute of  
Environmental Health Sciences

National Institutes of Health  
U.S. Department of Health and Human Services

## **Early Life Exposure to Outdoor Air Pollution and Respiratory Health, Ear Infections, and Eczema in Infants from the INMA Study**

Inmaculada Aguilera<sup>1,2</sup>, Marie Pedersen<sup>1,2,3</sup>, Raquel Garcia-Esteban<sup>1,2</sup>, Ferran Ballester<sup>2,4</sup>, Mikel Basterrechea<sup>2,5,6</sup>, Ana Esplugues<sup>2,4</sup>, Ana Fernández-Somoano<sup>2,7</sup>, Aitana Lertxundi<sup>6,8</sup>, Adonina Tardón<sup>2,8</sup>, and Jordi Sunyer<sup>1,2,9,10</sup>

<sup>1</sup> Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain

<sup>2</sup> Consortium for Research in Epidemiology and Public Health (CIBERESP), Spain

<sup>3</sup> National Institute of Health and Medical Research (INSERM), U823, Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, Grenoble, France

<sup>4</sup> Centre of Public Health Research (CSISP), Valencia, Spain

<sup>5</sup> Public Health Division of Gipuzkoa, Basque Country Government, Gipuzkoa, Spain

<sup>6</sup> BioDonostia Health Research Institute, Donostia, Spain.

<sup>7</sup> University of Oviedo, Oviedo, Spain

<sup>8</sup> Department of Public Health and Preventive Medicine, University of Basque Country, Bilbao, Spain

<sup>9</sup> Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain

<sup>10</sup> Pompeu Fabra University, Department of Experimental and Health Sciences, Barcelona, Spain.

### **Institution where the work was performed**

Centre for Research in Environmental Epidemiology  
Barcelona Biomedical Research Park  
Doctor Aiguader 88. 08003 Barcelona, Spain

### **Name and address of corresponding author**

Inmaculada Aguilera, PhD  
Centre for Research in Environmental Epidemiology  
Doctor Aiguader 88  
08003 Barcelona, Spain  
Phone: +34 932147300 ; Fax: +34 932147302  
E-mail: iaguilera@creal.cat

**Running title:** Air pollution and infant's health

**Keywords:** air pollution, children's health, ear infections, eczema, in utero exposure, respiratory infections.

**Acknowledgements:**

We thank all the participants for their generous collaboration. A full roster of the INMA project investigators can be found at [http://www.proyecto-inma.org/presentacion-inma/listado-investigadores/en\\_listado-investigadores.html](http://www.proyecto-inma.org/presentacion-inma/listado-investigadores/en_listado-investigadores.html). This study was funded by grants from Spanish Ministry of Health-Instituto de Salud Carlos III (Red INMA G03/176, CB06/02/0041, FIS-PI041436, FIS-PI081151, FIS-PI042018, FIS-PI09/02311, FIS-PI06/0867, FIS-PS09/00090, FIS-FEDER 03/1615, 04/1509, 04/1112, 04/1931, 05/1079, 05/1052, 06/1213, 07/0314, and 09/02647), Generalitat de Catalunya-CIRIT 1999SGR 00241, Conselleria de Sanitat Generalitat Valenciana, Universidad de Oviedo, Obra social Cajastur, Department of Health of the Basque Government (2005111093 and 2009111069), Provincial Government of Gipuzkoa (DFG06/004 and DFG08/001), and Fundación Roger Torné. Marie Pedersen holds a Juan de la Cierva post-doctoral fellowship awarded from the Spanish Ministry of Science and Innovation (JCI-2011-09479).

The authors declare they have no competing financial interests.

**Abbreviations**

GIS, Geographic Information Systems

INMA, Infancia y Medio Ambiente [Environment and Childhood]

IQR, interquartile range

LRTI, low respiratory tract infections

LUR, land use regression

NO<sub>2</sub>, nitrogen dioxide

## Abstract

**Background:** Prenatal and early life periods may be critical windows for harmful effects of air pollution on infant health.

**Objectives:** We studied the association of air pollution exposure during pregnancy and the first year of life with respiratory illnesses, ear infections, and eczema during the first 12-18 months of age in a Spanish birth cohort of 2,199 infants.

**Methods:** We obtained parentally-reported information on doctor-diagnosed lower respiratory tract infections (LRTI), and parental reports of wheezing, eczema, and ear infections. We estimated individual exposures to nitrogen dioxide (NO<sub>2</sub>) and benzene with temporally-adjusted land use regression models. We used log-binomial regression models and a combined random-effects meta-analysis to estimate the effects of air pollution exposure on health outcomes across the four study locations.

**Results:** A 10-μg/m<sup>3</sup> increase in average NO<sub>2</sub> during pregnancy was associated with LRTI (RR = 1.05; 95% CI: 0.98, 1.12) and ear infections (RR = 1.18; 95% CI: 0.98, 1.41). The RRs for an interquartile range (IQR) increase in NO<sub>2</sub> were 1.08 (95% CI: 0.97, 1.21) for LRTI and 1.31 (95% CI: 0.97, 1.76) for ear infections. Compared to NO<sub>2</sub>, the association for an IQR increase in average benzene exposure was similar for LRTI (RR = 1.06; 95% CI: 0.94, 1.19) and slightly lower for ear infections (RR = 1.17; 95% CI: 0.93, 1.46). Associations were slightly stronger among infants whose mothers spent more time at home during pregnancy. Air pollution exposure during the first year was highly correlated with prenatal exposure, thus we were unable to discern the relative importance of each exposure period.

**Conclusions:** Our findings support the hypothesis that early life exposure to ambient air pollution may increase the risk of upper and lower respiratory tract infections in infants.

## Introduction

Asthma, allergies, and respiratory infections are major health issues in childhood, and there is growing evidence that traffic-related air pollution is a risk factor for the development of asthmatic/allergic symptoms and respiratory infections (Braback and Forsberg 2009; Hoek et al. 2012). Ear infections are one of the leading causes of physician visits in young children as a major complication of a preceding upper respiratory infection, and have been moderately associated with ambient air pollution (Brauer et al. 2006; Macintyre et al. 2011; Zemek et al. 2010). The first years of life are thought to be the most important period for asthma and allergy development (Landau, 2006). As both the respiratory and immune systems are still under development, exposure to air pollution during this period may be critical for lasting adverse effects on respiratory health (Bateson and Schwartz 2008).

Most studies investigating the effects of outdoor air pollution on respiratory health, allergic symptoms, and ear infections during early childhood have focused on postnatal exposure, and only some of them have applied air pollution modelling techniques to obtain individual estimates of exposure (Brauer et al. 2002; Gehring et al. 2002; Karr et al. 2009; Macintyre et al. 2011; Morgenstern et al. 2007). There is increasing evidence from experimental and epidemiologic studies that the prenatal period is a critical window for harmful effects of toxic chemicals on respiratory health early in life and beyond (Pinkerton and Joad 2006). However, the effect of air pollution exposure during pregnancy on respiratory health and allergic responses early in life has been examined in a small number of studies that differed considerably with regard to exposure assessment and outcome definitions, which included physician records of asthma diagnosis (Clark et al. 2010), lung function measurements (Latzin et al. 2009), parental reports of respiratory infections and wheezing (Jedrychowski et al. 2005, 2011), and symptoms such as cough, sore throat, runny nose, or difficult breathing (Jedrychowski et al. 2005, 2011; Miller et al.

2004). Only two studies have assessed eczema (Jedrychowski et al. 2011; Miyake et al. 2010), and only one examined associations according to the time period of exposure during pregnancy (Esplugues et al. 2011).

Our aim was to investigate, in a Spanish population-based birth cohort, associations of exposure to outdoor air pollution during pregnancy and during the first year of life with respiratory illness, eczema, and ear infections during the first 12-18 months of life.

## **Methods**

### *Study population*

A population-based birth cohort was established between 2003 and 2008 in four Spanish regions (Asturias, Gipuzkoa, Sabadell, and Valencia) as part of the multicentre INMA-INfancia y Medio Ambiente [Environment and Childhood] project. Women were enrolled during the first routine prenatal care visit at the primary health care centre or hospital, provided that they fulfilled the inclusion criteria (age  $\geq 16$  years, wish to deliver in the reference hospital, no communication problems, singleton pregnancy, and no assisted conception) (Guxens et al. 2011). Follow-up interviews were performed during the third trimester of pregnancy, and for infants at 6 and 18 months in Asturias, at 14 months in Gipuzkoa, at 6 and 14 months in Sabadell, and at 12 months in Valencia. The Hospital Ethics Committees in the participating regions approved the study and all women provided written informed consent before participation.

### *Air pollution exposure assessment*

We developed area-specific land use regression (LUR) models of NO<sub>2</sub> and benzene to estimate residential-based exposures during pregnancy and the first year of life. Ambient levels of NO<sub>2</sub> and benzene were measured with Radiello passive samplers during several periods of one week each. Measurements taken in the different sampling campaigns were averaged to represent annual

mean levels in each study area. Potential predictor variables such as traffic indicators, surrounding land uses, topography, and population density were derived in the geographic information system ArcGIS 9.1 (ESRI, Redlands, CA, USA). Multiple linear regression models were built using a supervised forward stepwise procedure. Traffic-related variables, altitude, and land uses (urban, industrial, or agricultural) were the main predictor variables in the final LUR models. Models explained between 52 and 75% of the variability in measured NO<sub>2</sub> levels, and between 44 and 73% of the variability in measured benzene levels, depending on the study area. Residential addresses were geocoded using either commercial geocoding applications (in Asturias and Valencia) or mapping applications from the regional governments (in Gipuzkoa and Sabadell). LUR models were applied to predict outdoor levels of both pollutants at the residential addresses. For women and infants who changed their residential address during the study period, we calculated an average exposure estimate weighted by the time spent at each residence. We derived individual exposures to NO<sub>2</sub> and benzene during pregnancy by multiplying the LUR estimate by the ratio between the average concentrations measured at the fixed stations over the woman's pregnancy period and over the whole air pollution sampling period. We applied the same procedure to estimate exposures during each trimester of pregnancy and the first year of life. A description of the sampling campaigns, LUR models, and monitoring stations used for temporal adjustment in each of the cohorts is given in the Supplemental Material (Table S1).

#### *Definition of the outcomes*

Information on doctor-diagnosed LRTI, and symptomatic wheezing, eczema, and ear infections were obtained through interviewer-led questionnaires at each follow-up. Occurrence of a LRTI was defined as an affirmative answer to the question "Has a doctor told you that your son/daughter has had a chest infection?" An additional question referred to a specific doctor diagnosis of bronchitis, bronchiolitis, pneumonia, or pneumonitis. Parents were also asked if the

children had eczema, wheezing episodes, and ear infections from birth until follow-up. For these outcomes, parents were not specifically asked to report doctor-diagnosed conditions only. Due to funding constraints, the follow-up originally planned at 18 months in the Asturias cohort was actually performed at older ages (mean  $\pm$ SD = 29.1 $\pm$ 5.7 months), but in order to keep the adherence to the common study protocol mothers were specifically asked to refer to the period from 6 to 18 months of age when answering questions.

### *Covariates*

Potential confounding and effect modification variables (listed in Table 1) were obtained through questionnaires administered during the first and third trimesters of pregnancy, at birth, and during infants' follow-up. Cotinine levels were measured in a 24-hour sample of urine collected in the 3<sup>rd</sup> trimester of pregnancy using the Cotinine 175 Micro-Plate EIA Kit (Ora Sure Technologies, Inc.).

### *Statistical analysis*

We first performed area-specific analyses, using log-binomial regression models to estimate relative risks of outcomes during the first 12-18 months of life associated with an increase of 10  $\mu\text{g}/\text{m}^3$  and 1  $\mu\text{g}/\text{m}^3$  in  $\text{NO}_2$  and benzene exposures, respectively, for the entire pregnancy, each trimester, and the first year of life in each of the four study areas. We tested a subset of *a priori* selected covariates (defined as presented in Table 1) for inclusion in the models using backward selection, and we retained only those covariates associated with the outcome with  $P < 0.1$  or that led to a change in the relative risk estimate of at least 10% when removed from the model. In a second step, we performed a meta-analysis using random-effects models combining the relative risk estimates from each area, and estimated p-values for the Chi-Square test for heterogeneity. As a comparative analysis, we performed additional models using a biological approach, i.e. we included all the *a priori* selected covariates regardless of their p-value or effect on the exposure—



outcome association. We also estimated relative risks associated to an IQR increase in NO<sub>2</sub> and benzene exposure to be able to compare the effect of the two pollutants on health outcomes. Because LUR estimates of prenatal exposure were based on women's residential addresses, we performed a sensitivity analysis restricted to women who spent more time at home during pregnancy ( $\geq 15$  hours/day), assuming that this subset of participants is less affected by exposure misclassification (Nethery et al. 2009). Similarly, for postnatal exposures we estimated associations in the subset of infants that were not attending day care at the time of follow-up, under the assumption that they spent more time at or around their residence. Additional sensitivity analyses were conducted excluding infants from the Asturias cohort (given its particularities regarding infant's age at follow-up and the relevant contribution of industrial sources on air pollution levels), infants born preterm ( $< 37$  weeks of gestation), infants with low birth weight ( $< 2500$  g), and infants whose mothers smoked during pregnancy or after birth. Finally, we stratified by duration of breastfeeding (none,  $\leq 6$  months and  $> 6$  months), maternal fruits and vegetable intake during pregnancy (categorized using the median value), and maternal circulating vitamin D (categorized by tertiles) to examine potential effect modification by factors that may increase antioxidant and detoxification capacity, as suggested in a recent analysis of air pollution and neurodevelopment in the same cohort (Guxens et al. 2011). Statistical analyses were performed using Stata 10.1 (StataCorp, College Station, TX, USA).

## Results

From the 2,644 women enrolled in the study at the beginning of pregnancy, we obtained data on both air pollution exposure and health outcomes at 12-18 months for 83% ( $n=2,199$ ) of their infants (see Supplemental Material, Figure S1).

Characteristics of the study population are presented in Table 1. Most women were nulliparous (58%), nonsmokers during the entire pregnancy (82%) but exposed to passive smoking (62.5%), and had at least secondary education (77%). The percentage of maternal smoking increased in the postnatal follow-up (26%) but remained lower than paternal smoking (39%). The percentage of missing data among the covariates ranged from 0 to 5.4% (data not shown). Compared to the 2,199 mother-and-child pairs included in the study, the 445 non-participating women had lower educational level, higher prevalence of smoking during pregnancy, less prevalence of allergy history, and were less likely to be Spanish ( $P < 0.10$ ) (data not shown).

About 37% of the infants had a diagnosis of LRTI during the first 12-18 months of life (Table 2), with almost all (98%) classified as bronchitis or bronchiolitis. Thirty-eight percent of mothers reported that their child had one or more symptomatic episodes of wheezing, 33% reported one or more ear infections, and 22% reported at least one episode of eczema. All health outcomes were more likely to have been reported for children in the Asturias cohort, which also had the oldest infants' mean age at latest follow-up (mean $\pm$ SD = 29.2 $\pm$ 5.7 months compared with 13.6 $\pm$ 1.4 months for the three other cohorts combined). NO<sub>2</sub> exposure estimates were higher in the predominantly urban Valencia and Sabadell cohorts, while benzene levels were highest in Asturias, probably due to the additional contribution of industrial emissions (Figure 1). The correlation between NO<sub>2</sub> and benzene was considerably lower in Asturias (Spearman rho = 0.29) than in the other three areas (Spearman rho = 0.70 for all 3 combined). Levels of each pollutant were moderately to highly correlated between trimesters of pregnancy (Spearman rho = 0.62-0.79) and highly correlated between the entire prenatal period and the first year of life (Spearman rho = 0.94 for NO<sub>2</sub> and 0.89 for benzene).

Unadjusted and adjusted associations between exposure to air pollutants and development of health outcomes based on random-effects meta-analyses are shown in Table 3. RRs for a 10

$\mu\text{g}/\text{m}^3$  increase in  $\text{NO}_2$  exposure and 1  $\mu\text{g}/\text{m}^3$  increase in benzene exposure during the entire prenatal period suggest increased risk for LRTI and ear infections. The latter outcome is also positively associated with exposure to both pollutants during the first year of life. Relative risk estimates suggest a positive association between postnatal exposure to benzene and eczema but the rest of RRs for this outcome are close to unity. There are no substantial differences between crude and adjusted effect estimates for any of the two pollutants. Models adjusted for all of the potential covariates identified *a priori* produced slightly higher RRs with wider 95% CIs (results not shown). Results for an IQR increase in  $\text{NO}_2$  and benzene exposure during the entire pregnancy showed similar associations for both pollutants with LRTI (RR = 1.08; 95% CI: 0.97, 1.21 for  $\text{NO}_2$  and RR = 1.06; 95% CI: 0.94, 1.19 for benzene), whereas the association with ear infections was slightly higher for  $\text{NO}_2$  (RR = 1.31; 95% CI: 0.97, 1.76) than for benzene (RR = 1.17; 95% CI: 0.93, 1.46) (Supplemental Material, Table S2).

RRs for LRTI and ear infections in association with exposure to  $\text{NO}_2$  and benzene were slightly higher for exposure during the second trimester compared with other trimesters. Since most of the associations were adjusted by cotinine levels measured in urine at 32 weeks of pregnancy, which reflect very recent exposure to tobacco smoke, we ran additional models adjusted for questionnaire-based smoking variables instead, but associations remained essentially unchanged (data not shown).

With the exception of estimates for  $\text{NO}_2$  and ear infections, meta-analysis estimates did not show significant heterogeneity among the 4 study areas ( $P > 0.10$ , data not shown). Area-specific associations between  $\text{NO}_2$  and ear infections were positive for all time periods for all of the study areas except Sabadell (Supplemental Material, Table S3). In general, associations between ear infections and benzene were stronger in the two areas characterized by higher exposure levels, i.e. Asturias and Valencia. Because swimming pool attendance was particularly high in the Sabadell

cohort at 14 months of age (65% vs. 45% in the other cohorts), we stratified the analysis by this variable, but RRs for ear infections and NO<sub>2</sub> were comparable in both groups (results not shown). RRs for NO<sub>2</sub> (but not benzene) were slightly higher for infants whose mothers spent  $\geq 15$  hours/day at home during pregnancy compared with the cohort as a whole (1.08 vs. 1.05 for LRTI, 1.05 vs. 1.03 for wheezing, 1.04 vs. 1.00 for eczema, and 1.26 vs. 1.18 for ear infections) (Figure 2). Excluding the Asturias cohort did not show meaningful differences for any of the outcomes and resulted in wider confidence intervals for all the RRs for benzene exposure. Associations were essentially unchanged when preterm births and low birth weight children were excluded, as well as after excluding infants whose mothers smoked during pregnancy or after birth. Restricting analyses to infants who were not attending day care at the time of follow-up did not produce any clear trend (RRs associated with postnatal exposure to NO<sub>2</sub> decreased from 1.03 to 1.01 for LRTI, remained the same for wheezing, and increased from 1.02 to 1.04 for eczema and from 1.15 to 1.17 for ear infections, data not shown). Finally, we found no evidence of effect modification by maternal fruit and vegetable intakes, circulating vitamin D during pregnancy, or the duration of breastfeeding (Supplemental Material, Table S4).

## Discussion

In this prospective birth cohort study, we found associations with exposure to ambient air pollution during pregnancy and the first year of life with LRTI and ear infections during the first 12-18 months of life, although the high correlation between pre- and postnatal exposures did not allow us to identify whether either or both periods are driving the observed effects. We did not find clear or consistent evidence of associations between exposure to air pollution and eczema or wheezing symptoms. Associations were not substantially altered by adjusting for potential

confounders. In general, associations were slightly stronger for air pollution exposure during the second trimester of pregnancy compared with other trimesters.

We found similar patterns in the associations for NO<sub>2</sub> and benzene despite their moderate correlation. NO<sub>2</sub> is a widely used marker of traffic-related air pollution, whereas benzene levels can reflect both traffic and industrial activities, depending on the characteristics of the study area. Rather than a potential causative agent by itself, benzene may be acting as a surrogate for a mixture of predominantly traffic-driven pollutants, similar to NO<sub>2</sub>.

Differences in the proportions of children reported to have one or more episodes of each outcome may reflect differences in the length of follow-up among the four areas, which ranged from 12 to 18 months. All outcomes were more common in Asturias than in the other study areas, possibly because mothers reported on illnesses through 18 months when their children were 29 months of age on average, which could have led to over-reporting of outcomes. Exclusion infants from Asturias resulted in wider confidence intervals, particularly for RRs associated with benzene exposure, but associations were not modified to any great extent.

A major strength of our study is the prospective design beginning in early pregnancy, which allowed us to collect individual data on health outcomes and potential confounders using similar protocols in the four study areas. An additional strength is the use of temporally-adjusted LUR models to estimate individual exposures during specific time periods. However, despite their spatial accuracy, LUR estimates are still a proxy for personal exposure, which may be influenced by individual time-activity patterns (Nethery et al. 2009). Associations were slightly stronger when we limited analyses to infants whose mothers spent more time at home during pregnancy, both in the present study and in a previous analysis of birth outcomes (Estarlich et al. 2011), but we did not observe consistent changes in estimates after excluding infants attending day care at the time of follow-up.

The main limitation is that health data were obtained through questionnaires rather than clinical records. Recall bias and potential misclassification of the outcomes are possible, particularly in infants with less distinctive symptoms as compared to older children (Braback and Forsberg 2009).

Associations between NO<sub>2</sub> and LRTI, and to a lesser extent wheezing, are consistent with positive associations previously reported for respiratory infections and wheezing among 2-year-old infants in other European areas with similar NO<sub>2</sub> exposure levels (Brauer et al. 2002; Morgenstern et al. 2007). Wheezing at very young ages is highly associated with viral infections (Martínez et al. 1995), and many children who wheeze early in life have only a few episodes and do not wheeze at later ages. Nevertheless, even transient early wheeze is associated with asthma diagnosis later in childhood (Savenije et al. 2011).

Pre- and postnatal exposure to tobacco smoke is a risk factor for ear infections in early childhood, but evidence of an effect of ambient air pollution is inconclusive (Heinrich and Raghuyamshi 2004). Two recent studies reported modest but significant positive associations between short-term air pollution exposure and emergency room visits for ear infections (Zemek et al. 2010) and long-term exposure and physician diagnosis of otitis media (Macintyre et al. 2011). Similar to our study, a 10-μg/m<sup>3</sup> increase in postnatal exposure to NO<sub>2</sub> was associated with ear infections at 2 years of age in children from the Netherlands and Germany (Brauer et al. 2006).

Eczema is often the first manifestation of atopic disease during early childhood. Some of the factors that increase the risk of wheezing and respiratory infections in infants have less or no association with atopic eczema, which suggests differences in etiology between eczema and wheezing in infancy (Linneberg et al. 2006). Results from the few existing cohort studies on air pollution and eczema or allergic sensitization in young children are inconsistent, and comparisons are limited by differences in exposure assessment among studies (Brauer et al. 2002;

Jedrychowski et al. 2011; Miyake et al. 2010; Morgenstern et al. 2008; Nordling et al. 2008). A Dutch cohort that also used LUR models to estimate individual NO<sub>2</sub> exposures did not observe associations between doctor-diagnosed eczema at 2 years of age and NO<sub>2</sub> (OR=0.96; 95%CI: 0.85, 1.08 for a 10 µg/m<sup>3</sup> increase in NO<sub>2</sub>) (Brauer et al. 2002). The RRs close to unity for eczema in association with NO<sub>2</sub> exposure in our study are consistent with their findings.

Several studies have also estimated associations between prenatal exposure to air pollution and respiratory health outcomes early in life. Infants from a Polish cohort whose mothers were exposed to higher levels of polycyclic aromatic hydrocarbons during pregnancy (measured through 48-hour personal monitoring) had more frequent and longer episodes of several respiratory symptoms during the first year of life, including ear infections and wheezing (Jedrychowski et al. 2005). Pre- and postnatal exposures to several air pollutants, including NO<sub>2</sub>, have been positively associated with asthma in preschool age children (Clark et al. 2010). In a subset of 352 children from the present study, LUR estimates of NO<sub>2</sub> exposure levels during pregnancy and home outdoor measurements of NO<sub>2</sub> in the first year of life were not significantly associated with LRTI or with wheezing during the first year of life (Esplugues et al. 2011).

Air pollution exposures during pregnancy and first year of life tend to be highly correlated, which limits the interpretation of estimates from mutually adjusted models (Clark et al. 2010). In contrast with studies that characterize exposures based on measurements from nearest fixed monitoring stations (Mortimer et al., 2008), our exposure assessment approach emphasized spatial over temporal variation, which may have contributed to the very high correlations between pre- and post-natal exposures in our study. Results from the sensitivity analyses did not clarify the relative importance of each exposure period, because infants who did not attend day care were also more prone to be born from mothers who had spent more time at home during pregnancy. Therefore, it is not clear from our data which period (if any) was more relevant to health effects.

A pooled analysis of 12 studies on parental smoking and children's respiratory health suggested independent effects of pre- and postnatal tobacco smoke exposures (Pattenden et al. 2006).

However, associations between *in utero* exposure and respiratory outcomes in children appear to decline with age (Zlotkowska and Zejda 2005). From a biological perspective, both periods are likely to influence respiratory health given that lung development is a continuum of different phases that start in the embryogenesis and finish in early adolescence (Pinkerton and Joad 2006).

The exact mechanism whereby *in utero* exposure to ambient air pollution might increase the risk of postnatal respiratory infections is not known, but it has been hypothesized that air pollutants may affect the developing immune system which, could increase susceptibility to virally-induced infections after birth (Jedrychowski et al. 2005). We found slightly stronger associations with exposures during the second trimester of pregnancy, although differences must be interpreted with caution given the moderate to high correlations among trimester-specific exposures and the small differences in effect estimates. To our knowledge, apart from the study in the Valencia subsample of the INMA cohort (Esplugues et al. 2011), only one other cohort study of 232 asthmatic children assessed trimester-specific exposures during pregnancy in relation to allergic sensitization at 6-11 years old (Mortimer et al. 2008). The authors reported stronger associations for exposures during the first and second trimester of pregnancy, which suggests that alterations of developmental steps in this time period could lead to a shift of the Th1/Th2 balance in the immune system towards increased Th2 responses during later fetal development and after birth (Prescott et al. 2003). Further research on the effect of air pollution exposure during specific developmental periods *in utero* may contribute to identify the relevant mechanism(s) involved.



## Conclusion

This study supports the hypothesis that *in utero* or early exposure to ambient air pollution may affect the development of the respiratory system and increase the risk of upper and lower respiratory tract infections. The high correlation between exposures during pregnancy and first year of life did not allow us to disentangle associations with pre- versus post-natal exposures. Further study of this birth cohort is needed to determine whether exposure to air pollution during the first years of life is associated with the development of asthma at later ages.

## References

- Bateson TF, Schwartz J. 2008. Children's response to air pollutants. *J Toxicol Environ Health A* 71:238-243.
- Braback L, Forsberg B. 2009. Does traffic exhaust contribute to the development of asthma and allergic sensitization in children: findings from recent cohort studies. *Environ Health* 8:17.
- Brauer M, Hoek G, van Vliet P, Meliefste K, Fischer PH, Wijga A, et al. 2002. Air pollution from traffic and the development of respiratory infections and asthmatic and allergic symptoms in children. *Am J Respir Crit Care Med* 166:1092-1098.
- Brauer M, Gehring U, Brunekreef B, de Jongste J, Gerritsen J, Rovers M, et al. 2006. Traffic-related air pollution and otitis media. *Environ Health Perspect* 114:1414-18.
- Clark NA, Demers PA, Karr CJ, Koehoorn M, Lencar C, Tamburic L, et al. 2010. Effect of early life exposure to air pollution on development of childhood asthma. *Environ Health Perspect* 118:284-290.
- Esplugues A, Ballester F, Estarlich M, Llop S, Fuentes-Leonarte V, Mantilla E, et al. 2011. Outdoor, but not indoor, nitrogen dioxide exposure is associated with persistent cough during the first year of life. *Sci Total Environ* 409: 4667-4673.
- Estarlich M, Ballester F, Aguilera I, Fernandez-Somoano A, Lertxundi A, Llop S, et al. 2011. Residential exposure to outdoor air pollution during pregnancy and anthropometric measures at birth in a multicenter cohort in Spain. *Environ Health Perspect* 119:1333-1338.
- Gehring U, Cyrys J, Sedlmeir G, Brunekreef B, Bellander T, Fischer P, et al. 2002. Traffic-related air pollution and respiratory health during the first 2 yrs of life. *Eur Respir J* 19:690-698.
- Guxens M, Ballester F, Espada M, Fernandez MF, Grimalt JO, Ibarluzea J, et al. 2011. Cohort profile: The INMA-INfancia y Medio Ambiente-(Environment and Childhood) Project. *Int J Epidemiol*;doi:10.1093/ije/dyr054 [Online 5 April 2011]
- Guxens M, Aguilera I, Ballester F, Estarlich M, Fernandez-Somoano A, Lertxundi A, et al. 2012. Prenatal exposure to residential air pollution and infant mental development: modulation by antioxidants and detoxification factors. *Environ Health Perspect* 120:144-149.
- Heinrich J, Raghuyamshi VS. 2004. Air pollution and otitis media: a review of evidence from epidemiologic studies. *Curr Allergy Asthma Rep* 4:302-309.

- Hoek G, Pattenden S, Willers S, Antova T, Fabianova E, Braun-Fahrlander C, et al. PM10 and children's respiratory symptoms and lung function in the PATY study. *Eur Respir J*;doi: 10.1183/09031936.00002611 [Online 20 April 2012]
- Jedrychowski W, Galas A, Pac A, Flak E, Camman D, Rauh V, et al. 2005. Prenatal ambient air exposure to polycyclic aromatic hydrocarbons and the occurrence of respiratory symptoms over the first year of life. *Eur J Epidemiol* 20:775-782.
- Jedrychowski W, Perera F, Maugeri U, Mrozek-Budzyn D, Miller RL, Flak E, et al. 2011. Effects of prenatal and perinatal exposure to fine air pollutants and maternal fish consumption on the occurrence of infantile eczema. *Int Arch Allergy Immunol* 155:275-281.
- Karr CJ, Demers PA, Koehoorn MW, Lencar CC, Tamburic L, Brauer M. 2009. Influence of ambient air pollutant sources on clinical encounters for infant bronchiolitis. *Am J Respir Crit Care Med* 180:995-1001.
- Landau LI. Paediatric basis of adult lung disease. 2006. *Paediatr Respir Rev* 7 (Suppl 1):S251-254.
- Latzin P, Roosli M, Huss A, Kuehni CE, Frey U. 2009. Air pollution during pregnancy and lung function in newborns: a birth cohort study. *Eur Respir J* 33:594-603.
- Linneberg A, Simonsen JB, Petersen J, Stensballe LG, Benn CS. 2006. Differential effects of risk factors on infant wheeze and atopic dermatitis emphasize a different etiology. *J Allergy Clin Immunol* 117:184-189.
- Macintyre EA, Karr CJ, Koehoorn M, Demers PA, Tamburic L, Lencar C, et al. 2011. Residential air pollution and otitis media during the first two years of life. *Epidemiology* 22:81-89.
- Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. 1995. Asthma and wheezing in the first six years of life. The Group Health Medical Associates. *N Engl J Med* 332:133-138.
- Miller RL, Garfinkel R, Horton M, Camann D, Perera FP, Whyatt RM, et al. 2004. Polycyclic aromatic hydrocarbons, environmental tobacco smoke, and respiratory symptoms in an inner-city birth cohort. *Chest* 126:1071-1078.
- Miyake Y, Tanaka K, Fujiwara H, Mitani Y, Ikemi H, Sasaki S, et al. 2010. Residential proximity to main roads during pregnancy and the risk of allergic disorders in Japanese infants: the Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol* 21:22-28.

- Morgenstern V, Zutavern A, Cyrus J, Brockow I, Gehring U, Koletzko S, et al. 2007. Respiratory health and individual estimated exposure to traffic-related air pollutants in a cohort of young children. *Occup Environ Med* 64:8-16.
- Morgenstern V, Zutavern A, Cyrus J, Brockow I, Koletzko S, Kramer U, et al. 2008. Atopic diseases, allergic sensitization, and exposure to traffic-related air pollution in children. *Am J Respir Crit Care Med* 177:1331-1337.
- Mortimer K, Neugebauer R, Lurmann F, Alcorn S, Balmes J, Tager I. 2008. Air pollution and pulmonary function in asthmatic children: effects of prenatal and lifetime exposures. *Epidemiology* 19:550-557.
- Nethery E, Brauer M, Janssen P. 2009. Time-activity patterns of pregnant women and changes during the course of pregnancy. *J Expo Sci Environ Epidemiol* 19:317-324.
- Nordling E, Berglind N, Melen E, Emenius G, Hallberg J, Nyberg F, et al. 2008. Traffic-related air pollution and childhood respiratory symptoms, function and allergies. *Epidemiology* 19:401-408.
- Pattenden S, Antova T, Neuberger M, Nikiforov B, De Sario M, Grize L, et al. 2006. Parental smoking and children's respiratory health: independent effects of prenatal and postnatal exposure. *Tob Control* 15:294-301.
- Pinkerton KE, Joad JP. 2006. Influence of air pollution on respiratory health during perinatal development. *Clin Exp Pharmacol Physiol* 33:269-272.
- Prescott SL, King B, Strong TL, Holt PG. 2003. The value of perinatal immune responses in predicting allergic disease at 6 years of life. *Allergy* 58:1187-1194.
- Savenije OE, Granell R, Caudri D, Koppelman GH, Smit HA, Wijga A, et al. 2011. Comparison of childhood wheezing phenotypes in 2 birth cohorts: ALSPAC and PIAMA. *J Allergy Clin Immunol* 127:1505-1512.
- Zemek R, Szyszkowicz M, Rowe BH. 2010. Air pollution and emergency department visits for otitis media: a case-crossover study in Edmonton, Canada. *Environ Health Perspect* 118:1631-1636.
- Zlotkowska R, Zejda JE. 2005. Fetal and postnatal exposure to tobacco smoke and respiratory health in children. *Eur J Epidemiol* 20:719-727.

Table 1. Child and parental characteristics of the study population by study area

Characteristic	Asturias (n=434)	Gipuzkoa (n=540)	Sabadell (n=521)	Valencia (n=704)	Overall (n=2,199)
Male sex	51.8	49.9	52.0	52.8	51.7
Birth season					
Winter	21.9	29.4	26.1	25.7	25.9
Spring	19.4	28.4	30.5	18.9	24.1
Summer	30.4	24.3	23.2	21.3	24.3
Fall	28.3	17.8	20.2	34.1	25.7
Low birth weight	5.4	3.9	4.6	5.4	4.9
Preterm birth	5.1	3.4	3.1	5.3	4.3
Siblings at birth	38.9	42.1	41.5	44.6	42.2
Duration of breastfeeding (weeks)					
None	27.5	11.0	6.7	15.8	14.7
1-15	29.2	19.2	25.8	26.0	24.9
16-24	16.6	16.5	16.3	14.6	15.9
>24	26.7	53.3	51.2	43.6	44.6
Day care attendance	50.4	47.4	29.7	20.9	35.0
Non-Spanish mother	3.2	3.3	10.6	12.1	7.8
Pre-pregnancy BMI $\geq 25$ kg/m <sup>2</sup>	30.6	20.0	27.4	27.8	26.4

Table 1 (continued)

Characteristic	Asturias (n=434)	Gipuzkoa (n=540)	Sabadell (n=521)	Valencia (n=704)	Overall (n=2,199)
Maternal education					
Primary or less	17.7	13.5	27.4	31.5	23.4
Secondary	45.2	36.5	41.9	43.5	41.7
University	37.1	49.9	30.7	25.0	34.9
Maternal asthma	7.6	5.9	8.5	7.4	7.3
Paternal asthma	8.1	7.2	6.7	5.1	6.6
Maternal allergy	19.4	23.3	29.6	23.5	24.1
Paternal allergy	15.2	18.1	21.9	17.9	18.4
Maternal smoking during pregnancy	18.1	13.2	14.1	23.9	17.8
Secondhand smoke during pregnancy <sup>a</sup>					
No	50.7	42.1	36.9	26.8	37.5
1 source	35.6	43.2	42.7	39.0	40.3
2+ sources	13.7	14.7	20.4	34.2	22.2
Maternal postnatal smoking	26.1	19.4	25.4	31.3	26.0
Paternal postnatal smoking	33.2	31.5	41.5	45.8	38.9
Exposure to gas cooking during pregnancy	21.2	15.4	61.7	64.6	43.6
Mothers who spent ≥ 15 hr/day at home during pregnancy	91.7	42.5	52.5	62.8	60.9

Table 1 (continued)

Characteristic	Asturias (n=434)	Gipuzkoa (n=540)	Sabadell (n=521)	Valencia (n=704)	Overall (n=2,199)
Maternal fruits and vegetables intake during pregnancy <sup>b</sup>					
≤ 517.26 gr/day	47.7	48.3	47.2	56.0	50.4
> 517.26 gr/day	52.3	51.7	52.8	44.0	49.6
Maternal circulating vitamin D during pregnancy <sup>c</sup>					
≤ 20.89 ng/mL	39.1	36.6	38.2	23.0	33.2
20.90 - 30.20 ng/mL	36.1	35.8	28.9	34.2	33.7
> 30.20 ng/mL	24.8	27.7	32.9	42.7	33.1
Child's age at latest follow-up (months)	29.2 ± 5.7	14.3±1.2	14.5 ± 0.8	12.4 ± 1.1	16.59 ± 6.7
Maternal age at delivery (years)	33.0 ± 4.1	32.6±3.6	31.6 ± 4.2	31.3 ± 4.4	32.01 ± 4.2
Cotinine levels in urine at 32 weeks of gestation (ng/mL), log scale	2.2 ± 3.1	2.0±2.5	2.2 ± 2.8	3.0 ± 3.0	2.41 ± 2.9

Values are percentages for categorical variables and mean ± SD for continuous variables

<sup>a</sup> Sources of exposure to secondhand smoke reported during pregnancy: home, work, and other places

<sup>b</sup> Median cut-off value

<sup>c</sup> Tertile cut-off values

Table 2: Number and percentage of children reported to have had at least one episode of each health outcome during the first 12-18 months of age

	Asturias	Gipuzkoa	Sabadell	Valencia	Overall
Health outcome	(N = 434)	(N = 540)	(N = 521)	(N = 704)	(N = 2199)
Doctor-diagnosed					
LRTI	180 (44.7)	211 (39.1)	206 (40.9)	204 (29.0)	801 (37.2)
Wheezing	242 (59.3)	195 (36.1)	186 (37.7)	181 (25.7)	804 (37.5)
Eczema	148 (38.6)	88 (16.4)	107 (22.1)	117 (16.7)	460 (21.9)
Ear infections	173 (43.1)	194 (36.1)	174 (35.0)	172 (24.4)	713 (33.3)



Table 3. Association<sup>a</sup> between prenatal and postnatal exposure to outdoor NO<sub>2</sub> or benzene and doctor-diagnosed LRTI, wheezing, eczema, and ear infections during the first 12-18 months of age: results from random-effects meta-analysis across all four study areas.

Health outcome	Exposure period	NO <sub>2</sub> (per 10 µg/m <sup>3</sup> increase)					Benzene (per 1 µg/m <sup>3</sup> increase)				
		Crude		Adjusted			Crude		Adjusted		
		RR	(95% C.I.)	RR	(95% C.I.)	<i>p</i> -Value <sup>f</sup>	RR	(95% C.I.)	RR	(95% C.I.)	<i>p</i> -Value <sup>f</sup>
Doctor-diagnosed LRTI <sup>b</sup>	Entire prenatal	1.04	(0.97, 1.11)	1.05	(0.98, 1.12)	0.685	1.06	(0.97, 1.15)	1.05	(0.96, 1.14)	0.814
	First trimester	1.05	(0.99, 1.11)	1.06	(1.00, 1.12)	0.570	1.06	(1.00, 1.13)	1.06	(0.99, 1.13)	0.527
	Second trimester	1.07	(1.01, 1.13)	1.08	(1.02, 1.15)	0.336	1.09	(1.00, 1.19)	1.10	(1.01, 1.20)	0.279
	Third trimester	1.00	(0.94, 1.05)	1.00	(0.93, 1.07)	0.233	1.01	(0.89, 1.15)	0.99	(0.87, 1.12)	0.100
	First year of life	1.01	(0.95, 1.08)	1.03	(0.95, 1.11)	0.259	1.04	(0.95, 1.13)	1.02	(0.93, 1.11)	0.608
Wheezing <sup>c</sup>	Entire prenatal	1.02	(0.94, 1.10)	1.03	(0.96, 1.10)	0.631	1.03	(0.90, 1.17)	1.01	(0.94, 1.09)	0.396
	First trimester	1.02	(0.96, 1.07)	1.02	(0.96, 1.09)	0.918	0.99	(0.93, 1.05)	1.00	(0.94, 1.06)	0.929
	Second trimester	1.04	(0.96, 1.13)	1.05	(0.99, 1.11)	0.568	1.05	(0.92, 1.20)	1.02	(0.96, 1.09)	0.404
	Third trimester	0.99	(0.92, 1.07)	1.01	(0.94, 1.08)	0.299	0.99	(0.93, 1.05)	1.00	(0.93, 1.07)	0.531
	First year of life	1.03	(0.95, 1.11)	1.04	(0.98, 1.12)	0.486	0.98	(0.90, 1.08)	0.97	(0.90, 1.05)	0.597
Eczema <sup>d</sup>	Entire prenatal	1.00	(0.90, 1.10)	1.00	(0.91, 1.10)	0.819	1.03	(0.93, 1.14)	1.02	(0.90, 1.16)	0.355
	First trimester	0.97	(0.89, 1.06)	0.97	(0.89, 1.05)	0.486	0.94	(0.82, 1.09)	0.93	(0.81, 1.08)	0.171
	Second trimester	1.00	(0.92, 1.09)	1.01	(0.92, 1.11)	0.332	1.02	(0.89, 1.17)	1.02	(0.86, 1.22)	0.109
	Third trimester	1.01	(0.92, 1.10)	1.01	(0.92, 1.10)	0.815	1.07	(0.97, 1.19)	1.07	(0.97, 1.18)	0.363
	First year of life	1.01	(0.92, 1.11)	1.02	(0.92, 1.12)	0.924	1.10	(1.00, 1.22)	1.09	(0.98, 1.21)	0.690

Table 3 (continued)

Health outcome	Exposure period	NO <sub>2</sub> (per 10 µg/m <sup>3</sup> increase)					Benzene (per 1 µg/m <sup>3</sup> increase)				
		Crude		Adjusted			Crude		Adjusted		
		RR	(95% C.I.)	RR	(95% C.I.)	<i>p</i> -Value <sup>f</sup>	RR	(95% C.I.)	RR	(95% C.I.)	<i>p</i> -Value <sup>f</sup>
Ear infections <sup>c</sup>	Entire prenatal	1.14	(0.99, 1.33)	1.18	(0.98, 1.41)	0.002	1.13	(0.99, 1.28)	1.13	(0.95, 1.34)	0.121
	First trimester	1.07	(0.96, 1.20)	1.11	(0.99, 1.24)	0.048	1.06	(0.96, 1.17)	1.08	(1.02, 1.15)	0.726
	Second trimester	1.10	(0.95, 1.26)	1.16	(0.98, 1.37)	<0.001	1.07	(0.91, 1.26)	1.13	(1.00, 1.27)	0.120
	Third trimester	1.09	(0.96, 1.24)	1.12	(0.98, 1.29)	0.008	0.99	(0.85, 1.15)	1.02	(0.92, 1.13)	0.228
	First year of life	1.14	(1.02, 1.28)	1.15	(1.01, 1.31)	0.036	1.10	(1.10, 1.20)	1.08	(0.99, 1.18)	0.984

<sup>a</sup> Crude associations are adjusted for child’s sex and age at follow-up. Adjusted associations also include the following covariates depending on the study area:

<sup>b</sup> Day care attendance, siblings at birth, maternal asthma, parental allergy, maternal age at delivery, pre-pregnancy BMI, and cotinine levels in urine at 32 weeks of gestation (log- transformed).

<sup>c</sup> Duration of breastfeeding, day care attendance, siblings at birth, paternal allergy, parental asthma, maternal age at delivery, maternal education, pre-pregnancy BMI, maternal smoking during pregnancy, paternal post-natal smoking, and cotinine at 32 wks of gestation (log- transformed).

<sup>d</sup> Birth season, day care attendance, paternal asthma, maternal allergy, maternal age at delivery, pre-pregnancy BMI, maternal smoking during pregnancy, cotinine levels in urine at 32 weeks of gestation (log- transformed), and exposure to gas cooking during pregnancy.

<sup>e</sup> Birth season, day care attendance, siblings at birth, parental asthma and allergy, pre-pregnancy BMI, secondhand smoke exposure during pregnancy, and parental post-natal smoking.

<sup>f</sup> *p*-Value for the Chi-square test for heterogeneity

## Figure Legends

Figure 1. Distribution of exposures to outdoor NO<sub>2</sub> and benzene during pregnancy and first year of life by study area. Boxes extend from the 25th to the 75th percentile, horizontal bars represent the median, whiskers extend 1.5 times the length of the IQR above and below the 75th and 25th percentiles, respectively, and outliers are represented as points.

Figure 2. Adjusted associations from random-effects meta-analyses between exposure to outdoor NO<sub>2</sub> or benzene during pregnancy and LRTI, wheezing, eczema, and ear infections at 12-18 months of age in several subsamples of the cohort. Models were adjusted for the covariates indicated in Table 3.




Figure 1  
43x61mm (300 x 300 DPI)

